

CONNECTEDNESS OF PLANAR SELF-AFFINE SETS ASSOCIATED WITH NON-COLLINEAR DIGIT SETS

KING-SHUN LEUNG AND JUN JASON LUO

ABSTRACT. We study the connectedness of the planar self-affine sets $T(A, \mathcal{D})$ generated by an integer expanding matrix A with $|\det(A)| = 3$ and a non-collinear digit set $\mathcal{D} = \{0, v, kAv\}$ where $k \in \mathbb{Z} \setminus \{0\}$ and $v \in \mathbb{Z}^2$ such that $\{v, Av\}$ is linearly independent. By checking the characteristic polynomials of A case by case, we obtain a criterion concerning only k to determine the connectedness of $T(A, \mathcal{D})$.

1. Introduction

Let $M_n(\mathbb{Z})$ denote the set of $n \times n$ matrices with integer entries, let $A \in M_n(\mathbb{Z})$ be expanding, i.e., all eigenvalues of A have moduli strictly larger than 1. Assume $|\det A| = q$, and a finite set $\mathcal{D} = \{d_1, \dots, d_q\} \subset \mathbb{R}^n$ with cardinality q , we call it a *q-digit set*. It is well known that there exists a unique *self-affine set* $T := T(A, \mathcal{D})$ [12] satisfying:

$$T = A^{-1}(T + \mathcal{D}) = \left\{ \sum_{i=1}^{\infty} A^{-i} d_{j_i} : d_{j_i} \in \mathcal{D} \right\}.$$

T is called a *self-affine tile* if such T has a nonvoid interior.

The topological structure of $T(A, \mathcal{D})$, especially its connectedness, has attracted a lot of attentions in fractal geometry and number theory. It was asked by Gröchenig and Haas [5] that given an expanding integer matrix $A \in M_n(\mathbb{Z})$, whether there exists a digit set $\mathcal{D}$ such that $T(A, \mathcal{D})$ is a connected tile and they partially solved the question in $\mathbb{R}^2$. Hacon et al. [6] proved that any self-affine tile $T(A, \mathcal{D})$ with a 2-digit set is always pathwise connected. Lau and his collaborators ([8], [10], [11], [13]) systematically studied connectedness of the self-affine tiles generated by a kind of special digit sets of the form $\{0, 1, \dots, q-1\}v$ where $v \in \mathbb{Z}^n \setminus \{0\}$, which are called *consecutive collinear digit sets*. They observed a height reducing property (HRP) of the characteristic polynomial of A to determine the connectedness of $T(A, \mathcal{D})$, and conjectured that all monic expanding polynomials have the HRP, thus all the tiles generated by consecutive collinear digit sets are connected. Akiyama and Gjini [1] solved it up to degree 4. However it is still open for arbitrary degree. In the plane,

Date: August 21, 2012.

Key words and phrases. connectedness, self-affine set, digit set, $\mathcal{E}$ -connected.

The research is supported by STU Scientific Research Foundation for Talents.

the disk-likeness (i.e. homeomorphic to a closed unit disk) is an interesting topic, Bandt and Gelbrich [2], Bandt and Wang [3], and Leung and Lau [13] investigated the disk-likeness of self-affine tiles in terms of the neighborhoods of T . Deng and Lau [4], and Kirat [9] concerned themselves about a class of planar self-affine tiles generated by product digit sets.

With regard to other types of digit sets, there are few results about the connectedness of $T(A, \mathcal{D})$ generated by non-consecutive or non-collinear digit sets. In [14], by counting the neighborhoods of T , the authors made a first attempt to exploit the case of non-consecutive collinear digit sets with $|\det(A)| = 3$, and obtained a complete characterization for $T(A, \mathcal{D})$ to be connected or not. As a subsequent research, in the present paper, we will focus on the non-collinear digit sets. Precisely, we discuss the self-affine sets $T(A, \mathcal{D})$ generated by an expanding matrix A with $|\det(A)| = 3$ and a non-collinear digit set $\mathcal{D} = \{0, v, kAv\}$ for $k \in \mathbb{Z} \setminus \{0\}$ and $v \in \mathbb{R}^2$ such that $\{v, Av\}$ is linearly independent. By checking the characteristic polynomials of A case by case, we obtain a criterion concerning only k to determine the connectedness of $T(A, \mathcal{D})$.

Theorem 1.1. *Let A be a 2×2 integral expanding matrix with $|\det(A)| = 3$, and let $\mathcal{D} = \{0, v, kAv\}$ be a digit set where $k \in \mathbb{Z} \setminus \{0\}$ such that $\{v, Av\}$ is linearly independent. Then the self-affine set $T(A, \mathcal{D})$ is connected if and only if $k = \pm 1$.*

2. Preliminaries

In this section, we give some preparatory results of self-affine sets which will be used frequently in the paper. Define

$$\mathcal{E} = \{(d_i, d_j) : (T + d_i) \cap (T + d_j) \neq \emptyset, d_i, d_j \in \mathcal{D}\}.$$

We say that d_i and d_j are $\mathcal{E}$ -connected if there exists a finite sequence $\{d_{j_1}, \dots, d_{j_k}\} \subset \mathcal{D}$ such that $d_i = d_{j_1}, d_j = d_{j_k}$ and $(d_{j_l}, d_{j_{l+1}}) \in \mathcal{E}, 1 \leq l \leq k - 1$.

It is easy to check that $(d_i, d_j) \in \mathcal{E}$ if and only if

$$d_i - d_j = \sum_{k=1}^{\infty} A^{-k} v_k \quad \text{where} \quad v_k \in \Delta \mathcal{D} := \mathcal{D} - \mathcal{D}.$$

Then we get the following criterion of connectedness of a self-affine set.

Proposition 2.1. ([7], [10]) *A self-affine set T with a digit set $\mathcal{D}$ is connected if and only if any two $d_i, d_j \in \mathcal{D}$ are $\mathcal{E}$ -connected.*

In the following, we mainly consider the planar self-affine set $T(A, \mathcal{D})$ generated by a 2×2 integral expanding matrix A with $|\det A| = 3$ and a digit set $\mathcal{D} =$

$\{0, v, kAv\}$ such that $\{v, Av\}$ is linearly independent, where $k \in \mathbb{Z} \setminus \{0\}$. Denote the characteristic polynomial of A by $f(x) = x^2 + px + q$. Define α_i, β_i by

$$A^{-i}v = \alpha_i v + \beta_i Av, \quad i = 1, 2, \dots$$

Applying the Hamilton-Cayley theorem $f(A) = A^2 + pA + qI = 0$, it follows a lemma.

Lemma 2.2. ([13]) *Let α_i, β_i be defined as the above. Then $q\alpha_{i+2} + p\alpha_{i+1} + \alpha_i = 0$ and $q\beta_{i+2} + p\beta_{i+1} + \beta_i = 0$. Especially, $\alpha_1 = -p/q$, $\alpha_2 = (p^2 - q)/q^2$; $\beta_1 = -1/q$, $\beta_2 = p/q^2$. Moreover for $\Delta = p^2 - 4q \neq 0$, we have*

$$\alpha_i = \frac{q(y_1^{i+1} - y_2^{i+1})}{\Delta^{1/2}} \quad \text{and} \quad \beta_i = \frac{-(y_1^i - y_2^i)}{\Delta^{1/2}}$$

where $y_1 = \frac{-p+\Delta^{1/2}}{2q}$, $y_2 = \frac{-p-\Delta^{1/2}}{2q}$ are the two roots of $qx^2 + px + 1 = 0$.

Let

$$\tilde{\alpha} := \sum_{i=1}^{\infty} |\alpha_i|, \quad \tilde{\beta} := \sum_{i=1}^{\infty} |\beta_i|.$$

Corollary 2.3. *Assume $f(x) = x^2 + px + q$ and $g(x) = x^2 - px + q$ be the characteristic polynomials of expanding matrices A and B , respectively. Let $\alpha_i, \beta_i, \tilde{\alpha}, \tilde{\beta}$ for $f(x)$ be as before; let $\alpha'_i, \beta'_i, \tilde{\alpha}', \tilde{\beta}'$ be the corresponding terms for $g(x)$. Then*

$$\alpha'_{2j} = \alpha_{2j}, \quad \alpha'_{2j-1} = -\alpha_{2j-1}, \quad \beta'_{2j} = -\beta_{2j}, \quad \beta'_{2j-1} = \beta_{2j-1},$$

and hence $\tilde{\alpha} = \tilde{\alpha}'$, $\tilde{\beta} = \tilde{\beta}'$.

When $|\det A| = 3$, it is known by [2] that there are 10 eligible characteristic polynomials of A :

$$x^2 \pm 3; \quad x^2 \pm x + 3; \quad x^2 \pm 2x + 3; \quad x^2 \pm 3x + 3; \quad x^2 \pm x - 3.$$

Following [14], together with Corollary 2.3, we obtained the estimates or values of the corresponding $\tilde{\alpha}$ and $\tilde{\beta}$ as follows:

$$f(x) = x^2 \pm x + 3: \quad \tilde{\alpha} < 0.88, \quad \tilde{\beta} < 0.63; \tag{2.1}$$

$$f(x) = x^2 \pm 2x + 3: \quad \tilde{\alpha} < 1.17, \quad \tilde{\beta} < 0.73; \tag{2.2}$$

$$f(x) = x^2 \pm 3x + 3: \quad \tilde{\alpha} < 2.24, \quad \tilde{\beta} < 1.08; \tag{2.3}$$

$$f(x) = x^2 \pm x - 3: \quad \tilde{\alpha} = 2, \quad \tilde{\beta} = 1. \tag{2.4}$$

3. Main results

For a digit set $\mathcal{D} = \{0, v, kAv\}$, we denote by $\Delta\mathcal{D} = \{0, \pm v, \pm(kAv - v), \pm kAv\}$ the difference set. First we show the following simplest case according to the characteristic polynomials of A .

Theorem 3.1. *Let A be a 2×2 integral expanding matrix with characteristic polynomial $f(x) = x^2 \pm 3$ and $\mathcal{D} = \{0, v, kAv\}$ be a digit set where $k \in \mathbb{Z} \setminus \{0\}$ such that $\{v, Av\}$ is linearly independent. Then the self-affine set $T(A, \mathcal{D})$ is connected if and only if $k = \pm 1$.*

Proof. Since the case of $f(x) = x^2 - 3$ is more or less the same as that of $f(x) = x^2 + 3$, it suffices to show the last one. If $k = 1$, then $\Delta\mathcal{D} = \{0, \pm v, \pm(Av - v), \pm Av\}$. From $f(A) = A^2 + 3I = 0$, we have

$$I = -2A^{-2}(I + A^{-2})^{-1} = 2 \sum_{n=1}^{\infty} (-1)^n A^{-2n} \quad (3.1)$$

and

$$v = 2 \sum_{n=1}^{\infty} (-1)^n A^{-2n} v = \sum_{n=0}^{\infty} A^{-4n} (A^{-2}(-v) + A^{-3}(-Av) + A^{-4}v + A^{-5}(Av)). \quad (3.2)$$

Hence $v \in T - T$, or equivalently $T \cap (T + v) \neq \emptyset$. Moreover,

$$Av = \sum_{n=0}^{\infty} A^{-4n} (A^{-1}(-v) + A^{-2}(-Av) + A^{-3}v + A^{-4}(Av)) \quad (3.3)$$

which implies $T \cap (T + Av) \neq \emptyset$. Consequently, by Proposition 2.1, T is connected (see Figure 1(a)).




FIGURE 1. Two cases for $f(x) = x^2 + 3$ and $v = (1, 0)^t$.

If $k = -1$, then $\Delta\mathcal{D} = \{0, \pm v, \pm(Av + v), \pm Av\}$, and (3.1), (3.2), (3.3) still hold. Hence T is also connected.

If $|k| > 1$, let $k_i Av + l_i v \in \Delta \mathcal{D}$ for $i \geq 1$, then a point of $T - T$ can be written as

$$\begin{aligned}
& \sum_{i=1}^{\infty} A^{-i}(k_i Av + l_i v) = \sum_{i=1}^{\infty} A^{-2i}(k_{2i} Av + l_{2i} v) + \sum_{i=1}^{\infty} A^{-2i+1}(k_{2i-1} Av + l_{2i-1} v) \\
&= \sum_{i=1}^{\infty} \left(-\frac{1}{3}\right)^i (k_{2i} Av + l_{2i} v) + \sum_{i=1}^{\infty} \left(-\frac{1}{3}\right)^i (-3k_{2i-1} v + l_{2i-1} Av) \\
&= \left(k_1 + \sum_{i=1}^{\infty} \left(-\frac{1}{3}\right)^i (l_{2i} + k_{2i+1})\right) v + \left(\sum_{i=1}^{\infty} \left(-\frac{1}{3}\right)^i (l_{2i-1} + k_{2i})\right) Av \\
&:= Lv + KAv.
\end{aligned}$$

As $|l_i + k_{i+1}| \leq 1 + |k|$, it follows that $|K| \leq (1 + |k|) \sum_{i=1}^{\infty} \left(\frac{1}{3}\right)^i = (1 + |k|)/2 < |k|$. Hence $T \cap (T + kAv) = \emptyset$ and $(T + v) \cap (T + kAv) = \emptyset$, which imply that T is disconnected (see Figure 1(b)). $\square$

Theorem 3.2. *Let A be a 2×2 integral expanding matrix with characteristic polynomial $f(x) = x^2 + px \pm 3$ where $p > 0$, and let $\mathcal{D} = \{0, v, kAv\}$ be a digit set where $k \in \mathbb{Z} \setminus \{0\}$ such that $\{v, Av\}$ is linearly independent. Then the self-affine set $T(A, \mathcal{D})$ is connected if and only if $k = \pm 1$.*

Proof. For the cases of $f(x) = x^2 + px + 3$ with $0 < p < 3$, by using $0 = f(A) = f(A)(A - I) = A^3 + (p - 1)A^2 + (3 - p)A - 3I$, we obtain

$$I = \sum_{i=1}^{\infty} A^{-3i}((1 - p)A^2 - (3 - p)A + 2I). \quad (3.4)$$

Case 1. $f(x) = x^2 + x + 3$: For $k = 1$, then $\Delta \mathcal{D} = \{0, \pm v, \pm(Av - v), \pm Av\}$. By (3.4), $I = \sum_{i=1}^{\infty} A^{-3i}(-2A + 2I)$ and

$$v = \sum_{i=1}^{\infty} A^{-3i}(-2Av + 2v) = \sum_{i=0}^{\infty} A^{-3i}(A^{-2}(-v) + A^{-3}(v - Av) + A^{-4}(Av)). \quad (3.5)$$

Hence $T \cap (T + v) \neq \emptyset$. Moreover,

$$Av = \sum_{i=0}^{\infty} A^{-3i}(A^{-1}(-v) + A^{-2}(v - Av) + A^{-3}(Av)) \quad (3.6)$$

which implies $T \cap (T + Av) \neq \emptyset$. Consequently, T is connected (see Figure 2(a)).




FIGURE 2. Connected cases for $v = (1, 0)^t$ and $k = 1$.

For $k = -1$, then $\Delta\mathcal{D} = \{0, \pm v, \pm(Av + v), \pm Av\}$. From $f(A) = 0$, we deduce that $I = (-A - 2I)(A^2 + I)^{-1}$, which in turn gives

$$\begin{aligned}
 v &= -A^{-1}v - 2A^{-2}v + A^{-3}v + 2A^{-4}v - A^{-5}v - 2A^{-6}v + A^{-7}v + 2A^{-8}v - \dots \\
 &= A^{-2}(-Av - v) + A^{-3}(-Av) + A^{-4}(Av + v) + A^{-5}(Av) + A^{-6}(-Av - v) \\
 &\quad + A^{-7}(-Av) + A^{-8}(Av + v) + A^{-9}(Av) + \dots \\
 &\in T - T.
 \end{aligned}$$

Hence $T \cap (T + v) \neq \emptyset$. Multiplying the above expression by A , we have

$$\begin{aligned}
 Av &= A^{-1}(-Av - v) + A^{-2}(-Av) + A^{-3}(Av + v) + A^{-4}(Av) \\
 &\quad + A^{-5}(-Av - v) + A^{-6}(-Av) + A^{-7}(Av + v) + A^{-8}(Av) + \dots \\
 &\in T - T
 \end{aligned}$$

which implies $T \cap (T + Av) \neq \emptyset$. It follows that T is connected.

For $|k| > 1$. A point of $T - T$ can be written as

$$\sum_{i=1}^{\infty} A^{-i}(k_i Av + l_i v)$$

where $k_i Av + l_i v \in \Delta \mathcal{D}$ for $i \geq 1$. By using the relation $A^{-i}v = \alpha_i v + \beta_i Av$,

$$\begin{aligned} \sum_{i=1}^{\infty} A^{-i}(k_i Av + l_i v) &= \sum_{i=1}^{\infty} (k_i A^{-i+1}v + l_i A^{-i}v) \\ &= \sum_{i=1}^{\infty} k_i (\alpha_{i-1}v + \beta_{i-1}Av) + \sum_{i=1}^{\infty} l_i (\alpha_i v + \beta_i Av) \\ &= \left(k_1 + \sum_{i=1}^{\infty} (k_{i+1} + l_i) \alpha_i \right) v + \left(\sum_{i=1}^{\infty} (k_{i+1} + l_i) \beta_i \right) Av \\ &:= Lv + KAv. \end{aligned} \tag{3.7}$$

As $|l_i + k_{i+1}| \leq 1 + |k|$ and $\tilde{\beta} < 0.63$ (2.1), we conclude $|K| \leq 0.63(1 + |k|) < |k|$, which yields $T \cap (T + kAv) = \emptyset$ and $(T + v) \cap (T + kAv) = \emptyset$. Hence T is disconnected (see Figure 3(a)).

Case 2. $f(x) = x^2 + 2x + 3$: For $k = 1$. By (3.4), $I = \sum_{i=1}^{\infty} A^{-3i}(-A^2 - A + 2I)$ and

$$v = \sum_{i=1}^{\infty} A^{-3i}(-A^2 v - Av + 2v) = A^{-1}(-v) + \sum_{i=0}^{\infty} A^{-3i}(A^{-2}(-v) + A^{-3}v + A^{-4}(Av - v)).$$

Hence $T \cap (T + v) \neq \emptyset$. Moreover,

$$Av = A^{-1}(-Av) + \sum_{i=0}^{\infty} A^{-3i}(A^{-1}(-v) + A^{-2}v + A^{-3}(Av - v))$$

which implies $T \cap (T + Av) \neq \emptyset$. Consequently, T is connected (see Figure 2(b)).

For $k = -1$. We obtain from (3.4) that

$$\begin{aligned} v &= -A^{-1}v - A^{-2}v + 2A^{-3}v - A^{-4}v - A^{-5}v \\ &\quad + 2A^{-6}v - A^{-7}v - A^{-8}v + 2A^{-9}v + \dots \\ &= A^{-1}(-v) + A^{-2}(-v) + A^{-3}v + A^{-4}(Av) + A^{-5}(-Av - v) \\ &\quad + A^{-6}v + A^{-7}(Av) + A^{-8}(-Av - v) + \dots \\ &\in T - T, \end{aligned}$$

implying $T \cap (T + v) \neq \emptyset$. Multiplying the above expression by A , we have

$$\begin{aligned} Av + v &= A^{-1}(-v) + A^{-2}v + A^{-3}(Av) + A^{-4}(-Av - v) \\ &\quad + A^{-5}v + A^{-6}(Av) + A^{-7}(-Av - v) + \dots \\ &\in T - T, \end{aligned}$$

implying $(T + v) \cap (T - Av) \neq \emptyset$. Hence T is connected.




FIGURE 3. Disconnected cases for $v = (1, 0)^t$ and $k = 2$.

For $|k| > 1$. By (3.7) and $\tilde{\beta} < 0.73$ (2.2), we have $|K| \leq (1 + |k|)\tilde{\beta} < 0.73(1 + |k|)$. When $|k| \geq 3$, $|K| < 0.73(1 + |k|) < |k|$, which yields $T \cap (T + kAv) = \emptyset$ and $(T + v) \cap (T + kAv) = \emptyset$. Hence T is disconnected. When $k = 2$, suppose $(T + 2Av) \cap T \neq \emptyset$ or $(T + 2Av) \cap (T + v) \neq \emptyset$, i.e., $2Av + lv \in T - T$ for $l = 0$ or -1 . By (3.7), we obtain

$$(k_2 + l_1)\beta_1 = 2 - \sum_{i=2}^{\infty} (k_{i+1} + l_i)\beta_i \geq 2 - (1 + 2)(\tilde{\beta} - |\beta_1|) > 0.8 \quad (3.8)$$

where $\beta_1 = -1/3$. It follows that $l_1 = -1$, $k_2 = -2$. Then using $f(A) = 0$, we have

$$A(2Av + lv) - k_1Av - l_1v = (-4 + l - k_1)Av - (6 + l_1)v \in T - T.$$

It follows from $l_1 = -1$ that $k_1 = 0$ or 2 . Hence $-4 + l - k_1 \leq -4 + 0 + 0 = -4$, which contradicts the inequality $|K| \leq (1 + |k|)\tilde{\beta} < 3 \times 0.73 = 2.19$. So T is disconnected for $k = 2$ (see Figure 3(b)).

When $k = -2$, suppose $2Av + lv \in T - T$ where $l = 0$ or 1 . Similarly, it yields from (3.8) that $l_1 = -1$, $k_2 = -2$ and $k_1 = -2$ or 0 . If $k_1 = -2$, then

$(l-2)Av - 5v \in T - T$. Multiplying the expression by A and using $f(A) = 0$, we obtain

$$(l-2)A^2v - 5Av - (k_2Av + l_2v) = (1-2l)Av + (6-3l-l_2)v \in T - T$$

and $6-3l-l_2 \geq 6-3-1 = 2$. On the other hand, by (3.7) and (2.2), $-5.81 = -2-3\tilde{\alpha} \leq L \leq -2+3\tilde{\alpha} < 1.81$. This is ridiculous. If $k_1 = 0$, then $(l-4)Av - 5v \in T - T$ and $|l-4| \geq 3$ contracts $|l-4| = |K| < 2.19$. Therefore T is disconnected for $k = -2$.

Case 3. $f(x) = x^2 + 3x + 3$: For $k = 1$. From $(A-I)f(A) = 0$, we get $A^2 + A - I = 2(A+I)^{-1}$, which yields $I = -A^{-1} + A^{-2} + 2\sum_{i=3}^{\infty}(-1)^{i+1}A^{-i}$ and

$$\begin{aligned} v &= -A^{-1}v + A^{-2}v + 2\sum_{i=3}^{\infty}(-1)^{i+1}A^{-i}v \\ &= A^{-2}(v - Av) + A^{-3}v + \sum_{i=0}^{\infty}A^{-2i}(A^{-4}(Av - v) + A^{-5}(v - Av)). \end{aligned}$$

Hence $T \cap (T + v) \neq \emptyset$. Moreover,

$$Av = A^{-1}(v - Av) + A^{-2}v + \sum_{i=0}^{\infty}A^{-2i}(A^{-3}(Av - v) + A^{-4}(v - Av))$$

which implies $T \cap (T + Av) \neq \emptyset$. Consequently, T is connected (see Figure 2(c)).

For $k = -1$. From $f(A) = 0$ we get $A + 2I = -(A+I)^{-1}$. It follows that $I = -2A^{-1} - A^{-2} + A^{-3} - A^{-4} + A^{-5} + \dots$ and

$$\begin{aligned} v &= -2A^{-1}v - A^{-2}v + A^{-3}v - A^{-4}v + A^{-5}v + \dots \\ &= A^{-1}(-v) + A^{-2}(-Av - v) + A^{-3}v + A^{-4}(-v) + A^{-5}v + A^{-6}(-v) + \dots \\ &\in T - T, \end{aligned}$$

then $T \cap (T + v) \neq \emptyset$. Also we can deduce immediately that

$$\begin{aligned} Av + v &= A^{-1}(-Av - v) + A^{-2}v + A^{-3}(-v) + A^{-4}v + A^{-5}(-v) + \dots \\ &\in T - T, \end{aligned}$$

which yields $(T + v) \cap (T - Av) \neq \emptyset$. As a result, T is connected.

For $k > 1$. By (3.7) and (2.3), we have $|L| \leq k + (1+k)\tilde{\alpha} < 2.24 + 3.24k$. Suppose $kAv + lv \in T - T$ for $l = 0$ or -1 . Multiplying (3.7) by A and then subtracting $k_1Av + l_1v$ from both sides, we see that

$$(-3k + l - k_1)Av - (3k + l_1)v \in T - T. \quad (3.9)$$

Repeating the process, we obtain

$$(6k - 3l + 3k_1 - l_1 - k_2)Av + (9k - 3l + 3k_1 - l_2)v \in T - T. \quad (3.10)$$

Since $9k - 3l + 3k_1 - l_2 \geq 9k - 0 - 3k - 1 = 6k - 1 > 2.24 + 3.24k$, which exceeds the upper bound of $|L|$. It concludes that $T \cap (T + kAv) = \emptyset$ and $(T + v) \cap (T + kAv) = \emptyset$, that is, T is disconnected (see Figure 3(c)).

For $k \leq -3$, then $|L| \leq -k + (1 - k)\tilde{\alpha} < 2.24 - 3.24k$. It follows from (3.10) that $|9k - 3l + 3k_1 - l_2| \geq -9k - 3 + 3k - 1 = -6k - 4 > 2.24 - 3.24k$, which also exceeds the upper bound of $|L|$.

For $k = -2$, then $|K| < 3.24$ and $|L| < 8.72$. From (3.9), we have $6 + l - k_1 < 3.24$, hence $l = -1$ and $k_1 = 2$. From (3.10), we have $l_2 = -1$ and $3 + l_1 + k_2 < 3.24$, it follows that $l_1 = 0, k_2 = -2$, or $l_1 = 0, k_2 = 0$, or $l_1 = 1, k_2 = -2$.

When $l_1 = 0, k_2 = -2$. Multiplying (3.10) by A and then subtracting $k_3Av + l_3v$, we get

$$(-5 - k_3)Av + (3 - l_3)v \in T - T.$$

By $|5 + k_3| \leq 3.24$, it yields $k_3 = -2$. Repeating this process, we obtain

$$(12 - l_3 - k_4)Av + (9 - l_4)v \in T - T.$$

Hence we get a contradiction $|12 - l_3 - k_4| \geq 9 > 3.24$.

When $l_1 = 0, k_2 = 0$. Multiplying (3.10) by A and then subtracting $k_3Av + l_3v$, we get

$$(1 - k_3)Av + (9 - l_3)v \in T - T.$$

It yields $l_3 = 1$ and $k_3 = 0$ or 2 . Repeating this process, we obtain

$$(3k_3 + 5 - k_4)Av + (3k_3 - 3 - l_4)v \in T - T.$$

If $k_3 = 0$, then $(5 - k_4)Av + (-3 - l_4)v \in T - T$, and $k_4 = 2$. Finally we get $(-12 - l_4 - k_5)Av + (-9 - l_5)v \in T - T$ and a contradiction $|12 + l_4 + k_5| \geq 9 > 3.24$. If $k_3 = 2$, then $(11 - k_4)Av + (3 - l_4)v \in T - T$, and also $|11 - k_4| \geq 9 > 3.24$.

When $l_1 = 1, k_2 = -2$. By the same argument as above, we first get

$$(-2 - k_3)Av + (6 - l_3)v \in T - T.$$

It yields $k_3 = 0$. Repeating this process, we obtain $(12 - l_3 - k_4)Av + (6 - l_4)v \in T - T$ and $|12 - l_3 - k_4| \geq 9 > 3.24$ follows. Therefore T is disconnected for $|k| > 1$.

Case 4. $f(x) = x^2 + x - 3$: For $k = 1$. We deduce from $f(A) = 0$ that $I = (A^2 - I)^{-1}(-A + 2I)$, which yields $I = \sum_{i=0}^{\infty} A^{-2i}(-A^{-1} + 2A^{-2})$ and

$$\begin{aligned} v &= \sum_{i=0}^{\infty} A^{-2i}(-A^{-1} + 2A^{-2})v \\ &= A^{-2}(v - Av) + \sum_{i=1}^{\infty} A^{-2i}(A^{-1}(Av - v) + A^{-2}v). \end{aligned}$$

Hence $T \cap (T + v) \neq \emptyset$. Moreover,

$$Av = A^{-1}(v - Av) + \sum_{i=1}^{\infty} A^{-2i}((Av - v) + A^{-1}v)$$

which implies $T \cap (T + Av) \neq \emptyset$. Consequently, T is connected (see Figure 2(d)).

For $k = -1$. It follows from $v = A^{-1}(Av) \in T - T$ that $T \cap (T + v) \neq \emptyset$. Moreover, we can get $A + I = -I + (A - I)^{-1}$ from $f(A) = 0$. This implies

$$\begin{aligned} Av + v &= A^{-1}(-Av) + A^{-2}(Av) + A^{-3}(Av) + A^{-4}(Av) + \cdots \\ &\in T - T, \end{aligned}$$

that is, $(T + v) \cap (T - Av) \neq \emptyset$. Hence T is connected.

For $k > 1$. By (3.7) and (2.4), we have $|K| \leq (1 + k)\tilde{\beta} = 1 + k$ and $|L| \leq k + (1 + k)\tilde{\alpha} = 2 + 3k$. Suppose $kAv + lv \in T - T$ for $l = 0$ or -1 . Multiplying (3.7) by A and then subtracting $k_1Av + l_1v$ from both sides, we have

$$(-k + l - k_1)Av + (3k - l_1)v \in T - T.$$

Repeating the process, we obtain

$$(4k - l + k_1 - l_1 - k_2)Av + (-3k + 3l - 3k_1 - l_2)v \in T - T. \quad (3.11)$$

Note $4k - l + k_1 - l_1 - k_2 \geq 2k - 1$. When $k \geq 3$, $2k - 1 > k + 1$ which contradicts the upper bound of $|K|$, hence T is disconnected; when $k = 2$, it forces $l = 0$, $k_1 = -2$, $l_1 = 1$, $k_2 = 2$ and $3Av - l_2v \in T - T$, similarly which implies

$$(-3 - l_2 - k_3)Av + (9 - l_3)v \in T - T.$$

It is required that $|9 - l_3| \leq 8$, hence $l_3 = 1$. Furthermore, from $(-3 - l_2 - k_3)Av + 8v \in T - T$, we can deduce that

$$(11 + l_2 + k_3 - k_4)Av + (-9 - 3l_2 - 3k_2 - l_4)v \in T - T.$$

Since $11 + l_2 + k_3 - k_4 \geq 6 > 3$, we also get a contradiction, and T is disconnected. Consequently, T is disconnected for all $k > 1$ (see Figure 3(d)).

For $k < -1$, then $|K| \leq 1 - k$. From (3.11), it follows that $|K| \geq -4k + l - k_1 + l_1 + k_2 \geq -4k - 1 + k + 1 + k = -2k > 1 - k$, which is impossible. Therefore T is disconnected for $|k| > 1$. □

Theorem 3.3. *Let $B \in M_2(\mathbb{Z})$ be an expanding integral matrix with characteristic polynomial $g(x) = x^2 - px \pm 3$ where $p > 0$, and let $\mathcal{D} = \{0, v, kBv\}$ be a digit set where $k \in \mathbb{Z} \setminus \{0\}$ such that $\{v, Bv\}$ is linearly independent. Then the self-affine set $T(B, \mathcal{D})$ is connected if and only if $k = \pm 1$.*

Proof. The characteristic polynomial of A is $f(x) = x^2 + px + q$ if and only if that of $-A$ is $g(x) = x^2 - px + q$. Since only the radix expansions matter, we may assume $B = -A$. The proof for the disconnectedness of $T(B, \mathcal{D})$ when $|k| > 1$ can be adapted easily from the proof of Theorem 3.2 by applying Corollary 2.3. Let $\mathcal{D} = \{0, v, Av\} = \{0, v, -Bv\}$ and $\mathcal{D}' = \{0, v, -Av\} = \{0, v, Bv\}$. For $|k| = 1$, we deduce the connectedness of $T_1 := T(B, \mathcal{D})$ (respectively, $T'_1 = T(B, \mathcal{D}')$) from

that of $T = T(A, \mathcal{D})$ (respectively, $T' = T(A, \mathcal{D}')$). We only show the case of $f(x) = x^2 + x + 3$. In Case 1 of the proof of Theorem 3.2, from (3.5) we have

$$v = \sum_{i=0}^{\infty} (-B)^{-3i} (B^{-2}(-v) - B^{-3}(v + Bv) + B^{-4}(-Bv)) \in T_1 - T_1;$$

and from (3.6) we have

$$Bv = \sum_{i=0}^{\infty} (-B)^{-3i} (B^{-1}(-v) - B^{-2}(v + Bv) + B^{-3}(-Bv)) \in T_1 - T_1.$$

Hence T_1 is connected. Similarly, it can be verified that $v, Bv \in T'_1 - T'_1$ and T'_1 is also connected. $\square$

The above proof is indeed an application of the following more general result.

Theorem 3.4. *Let $A, B \in M_2(\mathbb{Z})$ be two expanding integral matrices with characteristic polynomials $f(x) = x^2 + px + q$ and $g(x) = x^2 - px + q$ respectively. Let v, w be two non-zero vectors such that the two sets $\{v, Av\}$ and $\{w, Bw\}$ are both linearly independent. Denote by L and M the lattices generated by $\{v, Av\}$ and $\{w, Bw\}$ respectively. Let $\mathcal{D} = \{c_i Av + d_i v \in L : i = 0, 1, \dots, |q| - 1\}$ and $\mathcal{D}' = \{-c_i Bw + d_i w \in M : i = 0, 1, \dots, |q| - 1\}$ be two digit sets. Then $T_1 = T(A, \mathcal{D})$ is connected if and only if $T_2 = T(B, \mathcal{D}')$ is connected.*

Proof. We may assume $B = -A$ and $w = -v$. Then $\mathcal{D} = \mathcal{D}'$ and $L = M$. Consider $a_i Av + b_i v = -a_i Bv + b_i v \in \Delta \mathcal{D} = \Delta \mathcal{D}'$, it follows that $a_i Av + b_i v = A^{-1}(c'_1 Av + d'_1 v) + A^{-2}(c'_2 Av + d'_2 v) + \dots \in T_1 - T_1$ if and only if $-a_i Bv + b_i v = B^{-1}(c'_1 Bv + d'_1 v) + B^{-2}(c'_2 Bv + d'_2 v) + \dots \in T_2 - T_2$. Hence $c_i Av + d_i v, c_j Av + d_j v \in \mathcal{D}$ are $\mathcal{E}$ -connected if and only if $-c_i Bv + d_i v, -c_j Bv + d_j v \in \mathcal{D}'$ are $\mathcal{E}$ -connected, then the theorem is proved by Proposition 2.1. $\square$

We can also deduce from Theorems 3.2 and 3.3 that

Corollary 3.5. *Let A be a 2×2 integral expanding matrix with characteristic polynomial $f(x) = x^2 + px + q$ where $|q| = 3$. Let $v \in \mathbb{R}^2$ such that $\{v, Av\}$ is linearly independent. Then the self-affine set $T(A, \mathcal{D})$ is connected for $\mathcal{D} = \{0, v, Av + v\}$ or $\{0, v, -Av + v\}$.*

Proof. Notice that the difference set $\Delta\{0, v, Av + v\} = \{0, \pm v, \pm Av, \pm(Av + v)\} = \Delta\{0, v, -Av\}$ and $\Delta\{0, v, -Av + v\} = \{0, \pm v, \pm Av, \pm(Av - v)\} = \Delta\{0, v, Av\}$. Hence the result follows from Proposition 2.1. $\square$

Remarks: The connectedness of self-affine sets or self-affine tiles is far from known extensively. Even for the planar case, there are still a lot of unsolved questions. The following may be some interesting topics related to the paper.

Q1. Can we characterize the connectedness of $T(A, \mathcal{D})$ with $|\det(A)| = 3$ and $\mathcal{D} = \{0, v, kAv + lv\}$?

Q2. For a two dimensional digit set $\mathcal{D} = \{0, v, \dots, (l-1)v, Av, \dots, kAv\}$ with $l+k = |\det(A)| > 3$, can we apply the same method to study the connectedness of $T(A, \mathcal{D})$?

Acknowledgments: The authors would like to thank Professor Ka-Sing Lau for suggesting a related question and advice on the work.

REFERENCES

- [1] S. Akiyama and N. Gjini, *Connectedness of number-theoretic tilings*, Discrete Math. Theoret. Computer Science 7 (2005), no. 1, 269-312.
- [2] C. Bandt and G. Gelbrich, *Classification of self-affine lattice tilings*, J. London Math. Soc. 50 (1994), 581-593.
- [3] C. Bandt and Y. Wang, *Disk-like self-affine tiles in $\mathbb{R}^2$* , Discrete Comput. Geom. 26 (2001), no.4, 591-601.
- [4] Q.R. Deng and K.S. Lau, *Connectedness of a class of planar self-affine tiles*, J. Math. Anal. Appl. 380 (2011) 493-500.
- [5] K. Gröchenig and A. Haas, *Self-similar lattice tilings*, J. Fourier Anal. Appl. 1 (1994), 131-170.
- [6] D. Hacon, N.C. Saldanha and J.J.P. Veerman, *Remarks on self-affine tilings*, Experiment. Math. 3 (1994), 317-327.
- [7] M. Hata, *On the structure of self-similar sets*, Japan J. Appl. Math. 2 (1985), no.2, 381-414.
- [8] X.G. He, I. Kirat and K.S. Lau, *Height reducing property of polynomials and self-affine tiles*, Geom. Dedicata (2011) 152:153-164.
- [9] I. Kirat, *Disk-like tiles and self-affine curves with non-collinear digits*, Math. Comp. 79 (2010), 1019-1045.
- [10] I. Kirat and K.S. Lau, *On the connectedness of self-affine tiles*, J. London Math. Soc. 6227 (2000), 291-304.
- [11] I. Kirat, K.S. Lau and H. Rao, *Expanding polynomials and connectedness of self-affine tiles*, Discrete Comput. Geom. 31 (2004), 275-286.
- [12] J.C. Lagarias and Y. Wang, *Self-affine tiles in $\mathbb{R}^n$* , Adv. Math. 121 (1996) 21-49.
- [13] K.S. Leung and K.S. Lau, *Disk-likeness of planar self-affine tiles*, Trans. Amer. Math. Soc. 359 (2007), 3337-3355.
- [14] K.S. Leung and J.J. Luo, *Connectedness of planar self-affine sets associated with non-consecutive collinear digit sets*, J. Math. Anal. Appl. 395 (2012) 208-217.

DEPARTMENT OF MATHEMATICS AND INFORMATION TECHNOLOGY, THE HONG KONG INSTITUTE OF EDUCATION, HONG KONG

E-mail address: ksleung@ied.edu.hk

DEPARTMENT OF MATHEMATICS, SHANTOU UNIVERSITY, SHANTOU 515063, CHINA

E-mail address: luojun@stu.edu.cn